FORM PTO-1390 US DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE ATTORNEY 'S DOCKET NUMBER MSKP039 TRANSMITTAL LETTER TO THE UNITED STATES U.S. APPLICATION NO (If known, see 37 CFR 1 5 DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371 INTERNATIONAL APPLICATION NO. INTERNATIONAL FILING DATE PCT/US99/10065 07 May 1999 08 May 1998 TITLE OF INVENTION COMPOSITIONS AND METHODS FOR ACTIVE VACCINATION EL556132244US APPLICANT(S) FOR DO/EO/US Agus, et al. Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information: 1. This is a FIRST submission of items concerning a filing under 35 US.C. 371. 2. This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 US.C. 371. 3. X This express request to begin national examination procedures (35 US.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C371(b) and PCT Articles 22 and 39(1). A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date A copy of the International Application as filed (35 U.S.C.371(c)(2)) is transmitted herewith (required only if not transmitted by the International Bureau). а has been transmitted by the International Bureau. b. is not required, as the application was filed in the United States Receiving Office (RO/US). A translation of the International Application into English (35 U.S.C. 371(c)(2)). Amendments to the claims of the International Aplication under PCT Article 19 (35 U.S.C. 371(c)(3)) are transmitted herewith (required only if not transmitted by the International Bureau) have been transmitted by the International Bureau. have not been made; however, the time limit for making such amendments has NOT expired. have not been made and will not be made A translation of the amendments to the claims under PCT Article 19 (35 U.S.C.371 (c)(3)). An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). A translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). rtems 11. to 16. below concern document(s) or information included: 11. An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included 13. A FIRST preliminary amendment. A SECOND or SUBSEQUENT preliminary amendment. 14. A substitute specification. 15. A change of power of attorney and/or address letter 16. Other items or information: Express Mail Cert. Exp. Mail No.: E L 556 13 22 44 US Date Mailed: 07 November Name:

528 Rec'd PCTPTO 07 NOV 2000

n9/ <i>(</i>	574975 PCT/U	S99/10065		MSKP039	
	ing fees are submitted:			CALCULATIONS	PTO USE ONLY
	FEE (37 CFR 1 492 (a)	. , . , , , ,			
nor international se	al preliminary examinati earch fee (37 CFR 1.445)	a)(2)) paid to HSPTO			
and International S	earch Report not prepare	d by the EPO or JPO	\$1070.00		
International prelin USPTO but Interna	rnational preliminary examination fee (37 CFR 1 482) not paid to PTO but International Search Report prepared by the EPO or JPO\$930.				
International prelin but international se	rnational preliminary examination fee (37 CFR 1 482) not paid to USPTO international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$790.				
	ninary examination fee (3 not satisfy provisions of F				
and all claims satis	ninary examination fee (3) affect provisions of PCT A	Article 33(1)-(4)	\$98.00		1
	R APPROPRIATE			\$ 710.00	
charge of \$130.0 nths from the earl	0 for furnishing the oath liest claimed priority date	or declaration later than e (37 CFR 1.492(e)).	20 30	\$	
LAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$	
tal claims	<u>∂</u> O -20 =	<u> </u>	x \$22.00	\$	
pendent claims	4 -3 =	11:1-1->	x \$82.00	\$ 82.00	
LTIPLE DEPEN	DENT CLAIM(S) (if app		+ \$270.00		
hotion of 1/2 for	filing by small entity, if	OF ABOVE CALCU		1100100	
t also be filed (N	Note 37 CFR 1.9, 1.27, 1.	28). Applicant class	ms small entity status. 4	\$ - 396.00	
:		See 37 CFR	UBTOTAL· =	\$ 396,00	1
essing fee of \$1	30.00 for furnishing the			\$	
ths from the ear	liest claimed priority date	e (37 CFR 1.492(f)).		Þ	
		TOTAL NATIO	NAL FEE =	\$ 396.00	
for recording the mpanied by an	e enclosed assignment (3 appropriate cover sheet (3	7 CFR 1.21(h)). The ass 37 CFR 3.28, 3 31). \$40	ignment must be .00 per property +	\$	
		TOTAL FEES E	NCLOSED =	\$ 396.00	
				Amount to be refunded:	\$
<u></u>				charged:	\$
b. Please char	the amount of \$3 or ge my Deposit Account is ecopy of this sheet is end	No ii	ne above fees is enclo		e above fees.
	issioner is hereby author nt to Deposit Account No				any
	appropriate time limit ust be filed and granted				ve (37 CFR
SEND ALL CORRESPO	ONDENCE TO:		m		lace 1
I HAARK TARKT HAARK			SIGNATIO	ma y 0)	ww.
	<u> </u>		Marina	T Largan	
02	1121		Marina	T. Larson	
 	: 1 4				
PATENT TRAD	EMARK OFFICE		32,038	ATION NUMBER	
			REGISTR	ATION NUMBER	

COMPOSITIONS AND METHODS FOR ACTIVE VACCINATION

This application relates to an active vaccine approach to the treatment of cancer and other diseases. The approach is applicable to a number of cancers and diseases, although a preferred embodiment provides an active vaccine for treatment of B cell Non-

Hodgkin's Lymphoma (NHL).

NHL is characterized by a clonal proliferation of malignant B cells. The treatment of NHL across a broad spectrum of patients remains a challenge, although numerous therapeutic approaches have been proposed and tried.

The most common therapeutic approach being used today is chemotherapy. While chemotherapy is effective for some period of time in most patients, a significant percentage of patients are not cured and experience a relapse.

Treatments have been proposed based on anti-idiotype therapy. In anti-idiotype therapy, a cell surface molecule which is expressed by malignant cells but not by normal cells is used to create patient-specific antibodies which are then administered to the patient. See, Miller, et al., *New Engl. J. Med.* 306: 517-522 (1982). Autologous patient-derived idiotype proteins have also been conjugated with keyhole limpet hemocyanin to produce a vaccine which has demonstrated efficacy and can elicit B and T cell immune responses. Kwak et al., *New Engl. J. Med.* 327: 1209-1215 (1992). Hybridoma-derived idiotype was co-cultured with patient-derived dendritic cells which acted as antigen presenters upon re-infusion into the patient and showed clinical efficacy. Hsu et al., *Nature Medicine* 2: 52-58 (1996). Idiotypic vaccines made in lipid-based carriers are disclosed in International Patent Publication WO98/14170.

Treatments have also been proposed using antibodies directed to CD20, a transmembrane protein that is expressed by both normal and malignant B-cells during parts of the B cell development cycle. Using single-dose infusions with anti-CD20 monoclonal antibodies, partial or minor tumor regressions were observed in 6 of 15 patients in a Phase I clinical study. Maloney et al., *Blood* 84: 2457-2466 (1994). In Phase II studies, 17 of 37 patients showed complete or partial remissions. In December 1997, the FDA approved the

5

25

30

5

first antibody-based therapy for NHL. Rituximab (Ritvaxan, IDEC/Genentech) is a chimeric human/murine antibody approved for the treatment of patients with relapsed or refractory low-grade or follicular CD20⁺ B cell NHL. Maloney et al., *Blood* 90: 2188-2195 (1997).

Combinations of chemotherapy and anti-CD20 therapy have been reported as having better therapeutic efficacy, with 11 of 11 patients showing complete or partial remission. Czuczman et al., Abstract 53, *Ann. Oncol.* 7, Supp. 1: 56 (1996).

While therapeutic regimens using anti-CD20 concepts are potentially effective, all of these therapies have the drawback of being passive therapies, i.e., they do not directly involve the immune system of the patient. Thus, these therapies may require the continued administration of the therapeutic agent for efficacy and do not provide any long-term protection against recurrence. In addition, the passive therapy is monoclonal in nature, therefore escape is possible. It would therefore be desirable to have an active therapy, that is a therapeutic agent which when administered to the patient stimulates an immune response against CD20 found in B-cells.

It is an object of the present invention to provide such a therapy. It is a further object of the invention to provide an active polyclonal therapy that is difficult to evade.

SUMMARY OF THE INVENTION

In accordance with the present invention, NHL is treated, not by administration of an anti-CD20 monoclonal antibody, but by the administration of CD20 itself, or an immunogenic fragment of the extracellular portion thereof, coupled to or administered with an antigenic carrier moiety such as keyhole limpet hemocyanin (KLH). This results in the stimulation of the production of polyclonal antibodies against CD20 (or an immunogenic fragment thereof) which has the affect of reducing the number of B-cells, including malignant B-cells. Thus, the invention provides an active vaccine. The same approach can be used for therapeutics for other diseases and conditions in which target cells possess a transmembrane protein, and is particularly applicable to those diseases and conditions for which administration of antibodies to transmembrane proteins or peptides (i.e., passive therapy) have been shown to provide therapeutic benefits, and especially in the

5

situations where the target is also capable of transducing or receiving a signal important for cell growth or function. This would include, for example, Her2/neu, VEGF receptor, epidermal growth factor receptor, the CD19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the P-glycoprotein, also known as the multidrug-resistance protein.

BRIEF DESCRIPTION OF THE FIGURES

Figs. 1A and B show ELISA results for formation of antibodies to human and mouse CD20 in vaccinated mice;

Figs. 2A and B shows results for binding of control B1 antibodies or antibodies in plasma from a mouse treated with human CD20-KLH conjugate with Raji B NHL cells;

Fig. 3 shows CP19⁺B cell levels in mice treated with human or mouse CD20-KLH conjugate;

Fig. 4 shows the domain structure of human Her2;

Fig. 5 shows the domain structure of human EGFR;

Figs. 6A-D shows the cross-reactivity of antibodies generated in response to human or mouse CD20 fragments;

Figs. 7A-D show the importance of carrier protein and adjuvant in generating an immune response;

Figs. 8A-D shows the immune response generated using different adjuvants; and

Figs. 9A-I shows CP19⁺B cell levels in mice treated with human or mouse CD20-KLH conjugate.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides an active vaccine therapy which can be used in the treatment of a variety of cancers and related conditions in which it is desirable to bring about the death of a target group of cells. Conventionally, immunotherapies targeting cells have sought to obtain a cellular immune response (T-cells that recognize the target cells), since a humoral immune response (antibodies that recognize the target cells) alone is not

30

25

20

25

30

deemed sufficient to achieve the desired result of cell death. The present invention departs from this conventional wisdom, and effectively utilizes a humoral immune response against the target cells to provide therapeutic benefit. The targets for therapy include cell surface proteins that when bound by a ligand signal to the cell. The vaccine induced antibody response will mimic ligand binding and cause similar signaling events which can imitate the process of programmed cell death (apoptosis) or halt the cell from growing or change the cancer cell's sensitivity to chemotherapy.

By way of example, the invention is suitably employed in the treatment of NHL and other B cell diseases such as chronic lymphocytic leukemia, auto-immune disorders and B-cell regulatory disorders. In accordance with this embodiment of the invention, a peptide antigen is prepared which contains at least an immunogenic portion of the extracellular domain of CD20 coupled to or administered with an antigenic carrier protein. The CD20 component of the peptide antigen may be syngeneic or it may be xenogeneic. Thus, for example, human patients may be treated with a peptide vaccine containing a human or a mouse CD20-fragment. There is evidence that strong immune responses can be elicited against xenogeneic proteins. Naftzger et al., Proc. Natl. Acad. Sci. (USA) 93: 14809-14814 (1996); International Patent Application PCT/US97/22669, filed December 10 1997, incorporated herein by reference. A suitable fragment is the 44 amino acid peptide spanning amino acids 136 to 179 of the sequence of mouse or human CD20. (Seq. ID Nos. 1 and 2) Other immunogenic fragments derived from the extracellular domain of CD20, or the entire CD20 molecule may also be used. Seq. ID. Nos. 3 and 4 shows the nucleic acid and amino acid sequences, respectively, of exon VI (the extracellular domain) of human CD20 as reported by Tedder et al., J. Immunol. 142: 2560-2568 (1989).

As used in the specification and claims hereof, an "immunogenic fragment" is a molecule which includes at least a portion of the extracellular domain of a transmembrane protein to direct and immunological response to that transmembrane protein when the immunogenic fragment is coupled to or administered with an antigenic carrier protein effective to break tolerance and administered with an adjuvant. It is not required that the immunogenic fragment alone be effective to stimulate an immune response, although such stimulation would not take a given fragment outside the scope of the present invention.

25

20

A preferred antigenic carrier protein is keyhole limpet hemocyanin which can be coupled to peptides using techniques described in Pierce Catalog Protocol. Other antigenic carrier proteins which can be used to break tolerance might be used in the invention include immunoglobulins, tuberculin, tetanus toxin and others well known in the art.

The peptide antigen containing the CD20 component and the antigenic carrier protein is formulated with a pharmaceutically acceptable adjuvant in a liquid carrier and administered to a patient suffering from NHL or another B cell disease. The composition will generally be administered by injection, for example, intramuscular, subcutaneous or intradermal injection, but might also be administered by way of a DNA vaccine (See US Patent No. 5,580,859, incorporated herein by reference) or a viral vaccine, or after mixing with antigen presenting cells (APC's) such as dendritic cells, *ex vivo*. Alternatively, the antigen may be administered without adjuvant by injection into a host prepared by prior or simultaneous injection of an immune adjuvant. Specific amounts to be administered to a patient can be determined by monitoring the titer of anti-CD20 antibodies developed by the patient, or by an average group of patients using well-known technology.

When a peptide of the extracellular domain of human or mouse CD20 is coupled to KLH and administered with an adjuvant to mice, antibodies which react with CD20 are found in plasma. (Figs. 1A and B) These antibodies bind to Raji cells, a human lymphoma cell line, indicating the ability to bind to a cell expressing CD20. (Figs. 2A and B). Moreover, the number of CD19⁺ B cells present in mice injected with either of the two CD20-KLH conjugates declines substantially (~30% decrease relative to controls). (Figs. 3 and 9). The assay used to quantitate B cell depletion detects CD19 which is also expressed on immature B cells that are CD20⁻. Thus, the 30% depletion actually underestimates the efficacy of the vaccine against CD20⁺ B cells.

Antibodies generated in mice after vaccination with human or mouse-derived CD20 fragments are specific for the peptides used, yet are capable of inducing immunity to the corresponding peptide from other species (Figs. 6A-D). Studies showed that in most instances the peptide, carrier protein and adjuvant are all needed for optimal response, although some responses were detected using less than all of the components. (Figs. 7A-D).

25

30

5

Several different adjuvants were also tested, and QS21 was found to be the most effective of those tested. (Fig. 8A-D).

While not intending to be bound by any particular mechanism, it is believed that the vaccines of the present invention are effective via at least two pathways. First, the generation of a humoral immune response to CD20 is effective to some extent to reduce the numbers of B cells bearing CD20 antigen in a manner consistent with normal immunological response to a target antigen. In addition, however, because CD20 has a signaling function, the binding of antibody to the CD20 moiety activates this signaling function to trigger apoptotic cell death. Such stimulation of apoptosis has been demonstrated to occur *in vitro* following passive treatments with a chimeric anti-CD20 antibody. Maloney et al., *Blood* 88 (Supp. 1): 637a (1996).

It is also possible that T cell mediated effector mechanisms are involved in the immune response. As evidence of this, we illustrate in Table 1 the mouse and human peptide sequences capable of binding to the corresponding mouse and human histocompatability antigens. This information was derived from a search of the NIH Bioinformatics and Molecular Analysis Section HLA Binding Predictions database using the mouse and human CD20 amino acid sequences. (Parker et al., *J. Immunol.* 152: 163 (1994)).

While the method of the invention is illustrated here using CD20 or CD20-derived peptides as the antigen to target B cells, the invention is not limited to this embodiment. Rather, the inventions encompasses the use of vaccine compositions comprising an immunogenic portions of the extracellular domain of transmembrane protein or peptide, particularly a transmembrane protein or peptide having signaling function, coupled to or administered with an antigenic protein and/or adjuvant to break tolerance.

A non-limiting example of another transmembrane protein which can be used in whole or in part in the method of the invention is Her-2/neu. The Her-2/neu oncogene is a receptor-like tyrosine kinase that is expressed on the cell surface of a significant portion of solid tumors. It has been shown that patients with early stage breast cancer have a high titer of antibodies to Her-2/neu. Disis et al., *J. Clin. Oncol.* 15: 3363-3367 (19967). The amino acid sequence and domain structure of human Her-2/neu are shown in Seq. OD. No. 5 and Fig. 4, and isolation and expression of the extracellular domain has been disclosed.

25

20

International Patent Publication No. WO 90/14357, which is incorporated herein by reference. There is clinical data showing efficacy of monoclonal antibodies against Her-2-neu in the treatment of patients with Her-2/neu⁺ tumors, and potential synergism with chemotherapy. Thus, in accordance with the present invention, a vaccine composition comprising at least an immunogenic portion of the extracellular domain of Her-2-neu (amino acids 22 to 652) coupled to or administered with an antigenic protein or peptide such a KLH can be used as a vaccine to provide the same therapeutic benefits using an active as opposed to a passive approach.

A further non-limiting example of a transmembrane protein which can be used in whole or in part in the method of the invention is epidermal growth factor receptor (EGFR). The amino acid sequence and domain structure of human EGFR are shown in Seq. ID. No. 6 and Fig. 5. There is significant data showing that antibodies to EGFR can have anti-tumor activity in breast and prostate cancer, as well as several head and neck tumors. Prewett et al., *J. Immunother. Emphasis Tumor Humoral* 19: 419-27 (1996). The mechanism by which antibody therapy against EGFR may be efficacious can be through the ability to down-regulate vascular endothelial growth factor production by tumor cells and thereby decrease angiogenesis. Petit et al., *Am. J. Pathol.* 151: 1523-30 (1997). In accordance with the present invention, a vaccine composition comprising at least an immunogenic portion of the extracellular domain of EGFR (amino acids 25 to 645) coupled to or administered with an antigenic protein or peptide such a KLH can be used as a vaccine to provide the same therapeutic benefits using an active as opposed to a passive approach. Preferred immunogenic peptides would be selected from regions not deleted in the various types of truncated EGFR mutants associated with some cancers.

A further non-limiting example of a transmembrane protein which can be used in whole or in part in the method of the invention is VEGF receptor. There are significant data showing that antibodies to VEGF receptor can inhibit angiogenesis and thereby halt tumor progression. In accordance with the present invention, a vaccine composition comprising at least an immunogenic portion of the extracellular domain of VEGF receptor coupled to or administered with an antigenic protein or peptide such a KLH can be used as a

vaccine to provide the same therapeutic benefits using an active as opposed to a passive approach.

Still a further non-limiting example of a transmembrane protein which can be used in whole or in part in the method of the invention is the IL-2 receptor. The IL-2 receptor is expressed on most T-cells malignancies, and there is a data showing that antibodies to the IL-2 receptor can be used in the treatment of T-cell malignancies and autoimmune disorders. In the present invention, a composition is made comprising at least an immunogenic portion of the extracellular domain of the IL-2 receptor (e.g., P55 or P75), coupled to or administered with an antigenic carrier protein or peptide such as KLH. and used as a vaccine.

The vaccine compositions of invention can be used alone or in combination (concurrently or sequentially) with drugs or chemotherapy agents that provide therapeutic benefit for the condition being treated. In the case of NHL, suitable chemotherapy agents which can be used in combination with the CD20 based vaccine include alkylating agents, anthrocyclines, cis-platinum, fludarabine, corticosteroids and vinca alkaloids. These same chemotherapy agents which might be used in combination with other vaccine compositions for other forms of cancer.

EXAMPLE 1

20

44 amino acid fragments of the extracellular domains of humans and murine CD20 (amino acids 136-179, Seq. ID Nos. 1 and 2) were synthesized using a solid-phase FMOC peptide synthesizer and coupled to KLH using the methodology described in the Pierce Catalog Protocol. The peptide coupled to KLH was then prepared for injection by formulation with QS-21 adjuvant. Balb/c mice were injected according to one of the following protocols on days 1, 8, 15, 22 and 50 of the experiment:

- A.
- Murine CD20 fragment-KLH with QS-21 adjuvant
- Human CD20 fragment-KLH with QS-21 adjuvant B.
- C. KLH with QS-21 adjuvant
- **OS-21** adjuvant D.

25

30

5

- E. P190 (irrelevant protein) coupled to KLH with QS-21 adjuvant
- F. B3A2 (irrelevant peptide) coupled to KLH with QS-21 adjuvant.

The animals were sacrificed on day 62 of the experiment.

Serum samples from the mice were diluted 1:200 and evaluated by BSA-blocked ELISA using goat-anti-mouse antibody conjugated to alkaline phosphatase for antibodies which bind to human CD20, mouse CD20 and KLH. As shown in Figs 1A and B, mice injected with human CD20 coupled to KLH (Fig. 1A) or mouse CD20 coupled to KLH (Fig. 1B) administration of xenogeneic antibody produced a significant polyclonal antibody response to both human and mouse CD20, while the response following administration of syngeneic antibody was principally limited to antibodies to the syngeneic form of CD20. Either xenogeneic or syngeneic peptide can therefore be used to generate an immune response.

To confirm the ability of the antibodies to bind to B cells, Raji cells (a form of human B-cell lymphoma that expresses CD20 on its surface) were blocked with human IgG, washed and then incubated for 30 minutes on ice with a 1:10 dilution of plasma from a mouse vaccinated with P-190-KLH control or huCD20-KLH. As a positive control, Raji cells were incubated with B1 antibody, or IgG2 as an isotypic negative control. After washing, the cells were incubated with goat-anti-mouse antibody, washed and fixed with 1% paraformaldehyde. Flow cytometry analysis was performed in a Becton-Dickinson FACScan. The results are shown in Figs 2A and 3B, wherein the shaded data set are the experimental data set and the outlined data set is the negative controls. As is apparent, there is a strong binding of mouse antibodies and Raji cells, comparable to that observed with B1 antibody.

25 <u>EXAMPLE 2</u>

To assess the number of B cells present in vaccinated mice, an evaluation was made of cells expressing CD19, a standard phenotypic marker for B cells. Spleens were harvested from the animals vaccinated in Example 1 and put into a single-cell suspension.

After counting the total number of cells, the cells were stained with FITC-labeled anti-mouse CD19 and the samples were analyzed by flow cytometry with a FACScan. 10,000 events

25

30

5

were collected. The percentage of CD19 positive cells minus the control gate was multiplied by the total number of cells to determine the number of CD19 positive cells in mice treated with the mouse and human CD20 peptide conjugates, and the P190 irrelevant peptide conjugate control.

As shown in Fig. 3, the absolute number of CD19 positive cells was significantly reduced in mice treated with either of the CD20 peptide conjugates. The level of CD19 positive cells is a reflection of the number of CD20 positive B cells, and the number of immature CD19⁺, CD20⁻ B cells in the samples. The absolute number of CD19⁺ B cells actually underestimates the therapeutic efficacy of the treatment for elimination of CD20⁺ B cells, however, since CD19 is expressed on B cell progenitor cells before expression of CD20.

EXAMPLE 3

Mice were injected five times over two months with one of four treatment protocols as follows:

human CD20 (44 aa fragment)-KLH plus QS1 human CD20 (44 aa fragment)-KLH

human CD20 (44 aa fragment) plus QS21

KLH plus QS21

Blood was collected on week 9 for analysis by ELISA. Sera from the vaccinated mice were diluted 1:200 and incubated on BSA blocked plates coated with msCD20, huCD20, P190 or KLH. Secondary goat anti-mouse antibody conjugated to alkaline phosphatase was added, and the color change of p-nitrophenyl phosphate substrate was measured at 405 nm. The results are summarized in Figs. 7A-D. In most instances the peptide, carrier protein and adjuvant are all needed for optimal response, although some responses were detected using less than all of the components.

EXAMPLE 4

Mice were vaccinated according to the schedule of Example 3 using one of four treatment protocols: human CD20 (44 aa fragment)-KLH plus QS21 adjuvant, mouse

5

CD20 (44 amino acid fragment)-KLH plus QS21, P190 (irrelevant protein)-KLH +QS21 and KLH and QS21 alone. Mouse serum samples were evaluated by ELISA for the presence of antibodies reactive with msCD20, huCD20, P190 and KLH. The results are shown in Figs. 6A-D. Antibodies generated in mice after vaccination with human or mouse-derived CD20 fragments are specific for the peptides used, yet are capable of inducing immunity to the corresponding peptide from other species.

EXAMPLE 5

Mice were vaccinated five times over two months with huCD20 fragment-KLH conjugate with no adjuvant or in combination with one of three adjuvants: QS21, BCG or Alum. Serum samples from the vaccinated mice were tested by ELISA. The results are summarized in Figs. 8A-D. QS21 was found to be the most effective of those tested.

EXAMPLE 6

To confirm the observations of Example 2, nucleated spleen cells were recovered by centrifugation in a density gradient from mice vaccinated with a CD20-KLH conjugate (human or mouse) in the presence of QS21 adjuvant. 1 X 10^6 cells from each mouse were incubated with 2 μ g of rat anti-mouse CD19 FITC-labeled antibody or with isotope-matched FITC labeled rat antibody. Cells were washed, fixed and analyzed with a Becton Dickinson FACScaliber cytometer. Figs 9A-C, D-F and G-I show the results for three exemplary mice of each vaccination group. The decrease in the peak reflecting levels of CD19 positive spleen cells in each of the mice is apparent.

,		-
	٥	2
_	c	5
	Ç	3

Score (T ₁₁₂ of Dissociation of Molecule Containing this Subsequence)	1600 48 21.2 1600 60 89.4 28.7
HLA Molecule	Kd Kd A_0201 Kd Kd A_0201 A_0201
Peptide Sequence	NFIRAHTPYI FIRAHTPYI FLKMeSLNFI HFLKmRRLEL IYDCePSNSS LIQTSKPYV ELIQtSKPYV
Species	human human human mouse mouse mouse

1

2

3 4

1

1

2

3

CLAIMS

- 1. A method for active vaccination against autologous cells expressing transmembrane proteins comprising administering to a patient a vaccine composition comprising at least an immunogenic portion of the extracellular domain of the transmembrane protein, or a xenogeneic homolog thereof, coupled to or administered with an carrier protein effective to break tolerance to the transmembrane protein and a pharmaceutically acceptable adjuvant.
- 2. The method of claim 1, wherein the transmembrane protein is selected from the group consisting of CD20, Her2-neu, VEGF receptor, epidermal growth factor receptor, the CD19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the P-glycoprotein.
 - 3. The method of claim 1, wherein the transmembrane protein is CD20.
- 4. The method of claim 1, wherein the vaccine composition comprises a peptide having the sequence given by Seq. ID No 1 or 2.
- 5. The method claim 1, wherein the carrier protein is keyhole limpet hemocyanin.
- 6. The method of claim 5, wherein the transmembrane protein is selected from the group consisting of CD20, Her2-neu, VEGF receptor, epidermal growth factor receptor, the CD19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the P-glycoprotein.
 - 7. The method of claim 5, wherein the transmembrane protein is CD20.

- 8. The method of claim 7, wherein the vaccine composition comprises a peptide having the sequence given by Seq. ID No 1 or 2.
 - 9. A method for active vaccination against B cells expressing CD20 comprising administering to a patient a vaccine composition comprising at least an immunogenic portion of the extracellular domain of CD20, or a xenogeneic homolog thereof, coupled to or administered with an carrier protein effective to break tolerance to the transmembrane protein and a pharmaceutically acceptable adjuvant.
 - 10. The method claim 9, wherein the carrier protein is keyhole limpet hemocyanin.
 - 11. The method of claim 9, wherein the vaccine composition comprises a peptide having the sequence given by Seq. ID No 1 or 2.
 - 12. A method for treatment of B cell non-Hodgkin's lymphoma, comprising administering to a patient suffering from B cell non-Hodgkin's lymphoma a vaccine composition comprising at least an immunogenic portion of the extracellular domain of CD20, or a xenogeneic homolog thereof, coupled to or administered with an carrier protein effective to break tolerance to the transmembrane protein and a pharmaceutically acceptable adjuvant.
 - 13. A vaccine composition comprising at least an immunogenic portion of the extracellular domain of the transmembrane protein, or a xenogeneic homolog thereof, coupled to or administered with an carrier protein effective to break tolerance to the transmembrane protein and a pharmaceutically acceptable adjuvant.
 - 14. The composition of claim 13, wherein the transmembrane protein is selected from the group consisting of CD20, Her2-neu, VEGF receptor, epidermal growth

3	factor receptor, the CD19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the
4	P-glycoprotein.
1 2	The composition of claim 13, wherein the transmembrane protein is CD20.
_	
1	16. The composition of claim 15, wherein the vaccine composition
2	comprises a peptide having the sequence given by Seq. ID No 1 or 2.
1 2 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	17. The composition of claim 13, wherein the carrier protein is keyhole limpet hemocyanin.
1	18. The composition of claim 17, wherein the transmembrane protein is
±2	selected from the group consisting of CD20, Her2-neu, VEGF receptor, epidermal growth
3	factor receptor, the CD19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the P-
3 4 1	glycoprotein. 19. The composition of claim 17, wherein the transmembrane protein is
2	CD20.
1	20. The composition of claim 19, wherein the vaccine composition
2	comprises a peptide having the sequence given by Seq. ID No 1 or 2.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:
A01N 37/18

(11) International Publication Number: WO 99/57981

(43) International Publication Date: 18 November 1999 (18.11.99)

(21) International Application Number: PCT/US99/10065 (81) Designated States: CA, JP, US, European patent (AT, BE, CH,

(22) International Filing Date: 7 May 1999 (07.05.99) PT, SE).

US

(71) Applicant (for all designated States except US): SLOAN-KETTERING INSTITUTE FOR CANCER RESEARCH [US/US]; 1275 York Avenue, New York, NY

8 May 1998 (08.05.98)

RESEARCH [US/US]; 1275 York Avenue, New York, NY 10021 (US).

(72) Inventors; and (75) Inventors/Applicants (for US only): AGUS, David, B.

(75) Inventors/Applicants (for US only): AGUS, David, B. [US/US]; 9 Pierrepont Street, Brooklyn, NY 11201 (US). SCHEINBERG, David [US/US]; 325 Central Park West, New York, NY 10025 (US). ROBERTS, Wendy [US/US]; 1233 York Avenue, New York, NY 10021 (US). ZELENETZ, Andrew, D. [US/US]; 31 Mohegan Road, Larchmont, NY 10538 (US).

(74) Agent: LARSON, Marina, T.; Oppedahl & Larson LLP, P.O. Box 5270, Frisco, CO 80443 (US). Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL,

(54) Title: COMPOSITIONS AND METHODS FOR ACTIVE VACCINATION


(57) Abstract

(30) Priority Data:

60/084,870

Non-Hodgkin's lymphoma (NHL) is treated, not by administration of an anti-CD20 monoclonal antibody, but by the administration of CD20 itself, or an immunogenic fragment of the extracellular portion thereof, coupled to or administered with an antigenic carrier moiety such as keyhole limpet hemocyanin (KLH). This results in the stimulation of the production of polyclonal antibodies against CD20 (or an immunogenic fragment thereof) which has the effect of reducing the number of B-cells, including malignant B-cells, and thus provides an active vaccine. The same approach can be used for therapeutics for other diseases and conditions in which target cells possess a transmembrane protein, and is particularly applicable to those diseases and conditions for which administration of antibodies to transmembrane proteins or peptides (i.e., passive therapy) have been shown to provide therapeutic benefits, and especially in the situations where the target is also capable of transducing or receiving a signal important for cell growth or function. This would include, for example, Her2/neu, VEGF receptor, epidermal growth factor receptor, the CD19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the P-glycoprotein, also known as the multidrug-resistance protein.










FIG. 1B



TOTAL EVENTS: 10000

	_,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		
MARKER	LEFT, RIGHT	EVENTS	% TOTAL
ALL	1, 9910	10000	100.00
M1	3, 17	7525	75.25
M2	1, 3	1201	12.01

FIG. 2A



TOTAL EVENTS: 10000

MARKER	LEFT, RIGHT	EVENTS_	% TOTAL
ALL	1, 9910	10000	100.00
M1	3, 12	7327	73.27
M2	1, 4	1454	1 4.54

FIG. 2B






FIG. 3











FIG. 6A

FIG. 6B







FIG. 6C

FIG. 6D







FIG. 7A

FIG. 7B






FIG. 7C

FIG. 7D




FIG. 8A




FIG. 8B




FIG. 8C





FIG. 8D






FIG. 9I

FILE NO. MSKP039 INVENTOR . Agus, et al

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My citizenship, residence and post office address are as listed below next to my name.

l believ which	ve I am the origina a patent is sought	al, first and [] sole on the invention	/[x] joint inventor of the subje entitled: <u>Compositions and M</u>	ect matter whic lethods for Acti	h is claimed and for ive Vaccination	
the spe	ecification of whicl	h				
(a)[]	is attached here	to.				
(p)[]	was filed on on		s Application Serial No.	,	_ and was amended	
(c) [X]	was described a and amended or	nd claimed in Inte	rnational Application No. <u>PC</u>	:T/US99/10065	5 filed on <u>May 7, 1999</u>	-
includir informa	ng the claims, as a ation which is mate	e reviewed and un amended by any a erial to the patenta	dedgment of Duty of Disclot derstood the content of the amendment referred to above ability of the subject matter of Regulations § 1.56(a).	above identified e. I acknowled	dae the duty to disclose	Э
365(c) insofar States acknow	of any PCT internation or PCT internation or PCT internation viedge the duty to the filing date of	ational application itter of each of the nal application in t disclose material	35 U.S.C. § 120 Inited States Code, § 120 of a designating the United State claims of this application is the manner provided by the finformation as defined in 37 dion and the national or PCT	tes of America, not disclosed i first paragraph CFR § 1.56 w	, listed below and, in the prior United of 35 U.S.C. § 112, I hich became available	>
(Applicati	ion Serial No.)	(Filing Date)	(Status)(patented,pending,aba	indoned)	(Patent No. if applicable)	
(Applicati	ion Serial No.)	(Filing Date)	(Status)(patented,pending,aba	indoned)	(Patent No. if applicable)	-
			Power of Attorney			
Nancy . Box 506	J. Parsons, PTO F 68, Alpine Bank C	Reg. No. 40,364 o enter, 2 nd Floor, 2	NO. 32,746, Marina T. Lars f the firm of OPPEDAHL & L 56 Dillon Ridge Rd., Dillon, G in the Patent and Trademan	ARSON LLP, h CO 80435 as a	having office at P.O.	
SEND	CORRESPONDENCE	ťo:	DIRECT TELEPHON OPPEDAHL & LARS((970) 468-6600		·	

PATENT TRADENARY OFFICE

(application number)



OPPEDAHL & LARSON

FILE NO. MSKP039 INVENTOR . Agus, et al

Claim for Priority

I hereby claim foreign priority benefits under 35 U.S.C. § 119 (a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below any foreign applications for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

APPLICATION	APPLICATION(S), FILED	WITHIN TWELVE MO	NTHS (6 MONTHS F	OR DESIGN) PRIC	OR TO SAID
COUNTRY	APPLICATION NO.	DATE OF FILING (day/month/year)	DATE OF ISSUE (day/month/year)	PRIORITY CLAIMED	CERTIFIED COPY ATTACHED
		· .		YES[]NO[]	YES[]NO[]
FOREIGN APPLICAT	ION(S), IF ANY, FILED MO	ORE THAN 12 MONTH	IS (6 MONTHS FOR I	DESIGN) PRIOR T	O SAID
COUNTRY	APPLICATION NO.	DATE OF FILING (day/month/year)	DATE OF ISSUE (day/month/year)		
hereby claim the b	enefit under 35 U.S.	Provisional App C § 119(e) of any		visional applica	tion(s) listed
60/084,870		08 Ma	y 1998		
(application number	7)	(filing d	ate)		

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

(filing date)

NAME OF SOLE OR FIRST INVENTOR	LAST NAME AGUS	FIRST NAME David	MIDDLE NAME B.
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE Brooklyn Beverly Hills	STATE OR COUNTRY OF RESIDENCE OF CA	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRESS STATES PIETEPONT Street	ss PA	CITY Brooklys PA Bevery Hills	STATE/COUNTRY ZIP CODE NY 10021 PA CA 90210
DATE III)	SIGNATURE	

[X] Signature for additional joint inventor attached. Number of Pages _1

[] Signature by Administrator(trix) or legal representative for deceased or incapacitated inventor. Number of Pages ___.

[] Signature for inventor who refuses to sign or cannot be reached by person authorized under 37 CFR § 1.47. Number of Pages ___.

08:33 10/30/5000

FILE NO. MSKP039 INVENTOR . Agus, et al

NAME OF SECOND INVENTOR	LAST NAME SCHEINBERG	FIRST NAME DAVID	MIDDLE NAME
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE NEW YORK	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRE 325 Central Park \		CITY New York	STATE/COUNTRY ZIP CODE NY 10025
DATE H	OO	SIGNATURE DOLLAR)
NAME OF THIRD INVENTOR	LAST NAME ROBERTS	FIRST NAME WENDY	MIDDLE NAME
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE NEW YORK N	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRESS 1233 York Avenue 303 East 71 St		CITY New York	STATE/COUNTRY ZIP CODE NY 10021
DATE ///DC)	SIGNATURE WILLIAM PHA	
NAME OF FOURTH INVENTOR	LAST NAME ZELENETZ	FIRST NAME ANDREW	MIDDLE NAME D.
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE LARCHMONT	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRESS 31 Mohegan Road		CITY Larchmont	STATE/COUNTRY ZIP CODE NY 10538
DATE 11/6/2000		SIGNATURE Auntolem	



OPPEDAHL & LARSON

FILE NO. MSKP039 INVENTOR . Agus, et al

COMBINED DECLARATION AND POWER OF ATTORNEY

MSKCC

As a below named inventor, I hereby declare that:

My citizenship, residence and post office address are as listed below next to my name.

I believe I am the original, first and [] sole/[x] joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled: Compositions and Methods for Active Vaccination

			Davies of Atlantas		
(Applicat	ion Serial No.)	(Filing Date)	(Status)(patented,pending,at	pandoned)	(Patent No. if applicable)
(Applicat	ion Serial No.)	(Filing Date)	(Status)(patented,pending,al	pandoned)	(Patent No. if applicable)
365(c) insofar States acknow	of any PCT internations the subject matter or PCT international vedge the duty to dien the filing date of the subject of the subject of the filing date of the subject of the filing date of the filing dat	onal application or of each of the application in the sclose material i	35 U.S.C. § 120 nited States Code, § 120 or designating the United States claims of this application is ne manner provided by the information as defined in 3 ion and the national or PCT	ates of America, I s not disclosed in first paragraph o 7 CFR § 1.56 whi f international filli	isted below and, the prior United f 35 U.S.C. § 112, I ich became available ng date of this
includir informa	ng the claims, as am ation which is materi	eviewed and uncomended by any a al to the patenta	degment of Duty of Discleterstood the content of the mendment referred to about bility of the subject matter (Regulations § 1.56(a).	above identified: /e. I acknowledg	e the duty to disclose
(c) [X]	was described and and amended on	claimed in Inter	national Application No. <u>P</u> (<u>CT/US99/10065</u> f	iled on <u>May 7, 1999</u>
(p)[]	was filed on	as	s Application Serial No	,	and was amended
(a)[]	is attached hereto.		•		
the spe	ecification of which			•	

Power of Attorney

I hereby appoint Carl Oppedahl, PTO Reg. No. 32,746, Marina T. Larson, PTO Reg. No. 32,038, and Nancy J. Parsons, PTO Reg. No. 40,364 of the firm of OPPEDAHL & LARSON LLP, having office at P.O. Box 5068, Alpine Bank Center, 2nd Floor, 256 Dillon Ridge Rd., Dillon, CO 80435 as attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith.

SEND CORRESPONDENCE TO:



DIRECT TELEPHONE CALLS TO: OPPEDAHL & LARSON LLP (970) 468-6600

PAYENT TRACEMEN OFFICE

(application number)



FILE NO. MSKP039 INVENTOR . Agus, et al

Claim for Priority

I hereby claim foreign priority benefits under 35 U.S.C. § 119 (a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below any foreign applications for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

EARLIEST FOREIGN APPLICATION(S), FILED WITHIN TWELVE MONTHS (6 MONTHS FOR DESIGN) PRIOR TO SAID APPLICATION						
COUNTRY	APPLICATION NO.	DATE OF FILING (day/month/year)	DATE OF ISSUE (day/month/year)	PRIORITY CLAIMED	CERTIFIED COPY ATTACHED	
				YES[]NO[]	YES[]NO[]	
FOREIGN APPLICAT APPLICATION	FOREIGN APPLICATION(S), IF ANY, FILED MORE THAN 12 MONTHS (6 MONTHS FOR DESIGN) PRIOR TO SAID APPLICATION					
COUNTRY	APPLICATION NO.	DATE OF FILING (day/month/year)	DATE OF ISSUE (day/month/year)			
Provisional Application hereby claim the benefit under 35 U.S.C § 119(e) of any United States provisional application(s) listed below.						
60/084,870	60/084,870 08 May 1998					
(application number) (filing date)						

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

(filing date)

NAME OF SOLE OR FIRST INVENTOR	LAST NAME AGUS	FIRST NAME David	MIDDLE NAME B.
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE Brooklyn	STATE OR COUNTRY OF RESIDENCE	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRESS 9 Pierrepont Street		CITY Brooklyn	STATE/COUNTRY ZIP CODE NY 10021
DATE		SIGNATURE	

[X] Signature for additional joint inventor attached. Number of Pages 1

[] Signature by Administrator(trix) or legal representative for deceased or incapacitated inventor. Number of Pages

[] Signature for inventor who refuses to sign or cannot be reached by person authorized under 37 CFR § 1.47. Number of Pages ___.



OPPEDAHL & LARSON

FILE NO. MSKP039 INVENTOR . Agus, et al

	T		
NAME OF SECOND INVENTOR	LAST NAME SCHEINBERG	FIRST NAME DAVID	MIDDLE NAME
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE NEW YORK	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRESS 325 Central Park West		CITY New York	STATE/COUNTRY ZIP CODE NY 10025
DATE		SIGNATURE	
NAME OF THIRD INVENTOR	LAST NAME ROBERTS	FIRST NAME WENDY	MIDDLE NAME
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE NEW YORK	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRESS 1233 York Avenue		CITY New York	STATE/COUNTRY ZIP CODE NY 10021
DATE		SIGNATURE	
NAME OF FOURTH INVENTOR	LAST.NAME ZELENETZ	FIRST NAME ANDREW	MIDDLE NAME D.
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE LARCHMONT	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRESS 31 Mohegan Road		CITY Larchmont	STATE/COUNTRY ZIP CODE NY 10538
DATE		SIGNATURE	

SEQUENCE LISTING

```
<110> Agus, David B.
         Scheinberg, David
         Zelenetz, Andrew D.
         Roberts, Wendy
  <120> Compositions and Methods for Active Vaccination
  <130> MSKP039US
  <140>
  <141>
  <150> 60/084,870
  <151> 1998-05-08
  <160> 6
 <170> PatentIn Ver. 2.0
 <210> 1
 <211> 44
 <212> PRT
 <213> HUMAN
 <220>
 <223> human CD20 fragment
 <400> 1
 Lys Ile Ser His Phe Leu Lys Met Glu Ser Leu Asn Phe Ile Arg Ala
                 5
                                      10
His Thr Pro Tyr Ile Asn Ile Tyr Asn Cys Glu Pro Ala Asn Pro Ser
             20
                                25
Glu Lys Asn Ser Pro Ser Thr Gln Tyr Cys Tyr Ser
         35
                             40
<210> 2
<211> 44
<212> PRT
<213> Murine
<220>
```

<223> murine CD20 fragment

```
<400> 2
 Thr Leu Ser His Phe Lou Lys Met Arg Arg Leu Glu Leu Ile Gln Thr
                                       10
 Ser Lys Pro Tyr val App fle Tyr App Cys Glu Pro Ser App Ser Ser
                                   25
 Glu Lys Asn Ser Pro Ser Thr Gln Tyr Cys Asn Ser
                               40
          35
 <210> 3
 <211> 79
 <212> PRT
 <213> HUMAN
 <220>
 <223> exon VI of numan CD20 gene
 <400> 3
 Val Lys Gly Lys Met Ile Met Asn Ser Lou Ser Leu Phe Ala Ala Ile
                                      10
                   5
  1
 Ser Gly Met Ile Leu Ser Ile Met Asp Ile Leu Asn Ile Lys Ile Ser
                                  25
              20
His Phc Lou Lys His Glu Ser Leu Asn Phe Ile Arg Ala His Thr Pro
                              40
          35
Tyr Ile Asn Ile Tyr Asn Cys Glu Pro Ala Asn Pro Ser Glu Lys Asn
                                              60
                         55
Ser Pro Ser Thr Glm Tyr Cys Tyr Ser Ile Glm Ser Leu Phe Leu
                     70
 65
<210> 4
<211> 237
<212> DNA
<213> HUMAN
<220>
<223> exon VI of human CD20 gene
<400> 4
gucaaaggaa aaatgataat gaattcattg agcototttg otgocattto tggaatgatt 60
ctttcaatca tggacatast taatattaaa atttcccatt ttttaaaaaat ggagagtctg 120
```

aattttatta gageteasas accatatatt aacatataca actgtgaace agetaateee 180

totgagaaaa actooccato tacccaatac tgttacagca tacaatotot gttottg 237

<210> 5 <211> 1255 <212> PRT <213> HUMAN <220> <223> human Her2 <400> 5 Met Glu Leu Ala Ala Leu Cys Arg Trp Gly Leu Leu Ala Leu Leu 10 Pro Pro Gly Ala Ala Ser Thr Gln Val Cys Thr Gly Thr Asp Met Lys 20 25 Leu Arg Leu Pro Ala Ser Pro Glu Thr His Leu Asp Met Leu Arg His 35 40 Lou Tyr Gln Gly Cys Gln Val Val Gln Gly Asn Lou Gl. Lou Thr Tyr 50 55 Leu Pro Thr Asn Ala Ser Leu Ser Phe Leu Gln Asp Ile Gln Glu Val 65 70 75 Gln Gly Tyr Val Leu Ile Ala His Asn Gln Val Arg Gln Val Pro Leu 85 90 Gln Arg Leu Arg Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Asn Tyr 105 Ala Leu Ala Val Leu Asp Asn Gly Asp Pro Leu Asn Asn Thr Thr Pro 115 120 Val Thr Gly Ala Ser Pro Gly Gly Leu Arg Glu Leu Gln Leu Arg Ser 130 135 Leu Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Gln Arg Asn Pro Gln 145 150 Leu Cys Tyr Gln Asp Thr Ile Leu Trp Lys Asp Ile Phe His Lys Asn 135 170 Asn Glm Leu Ala Leu Thr Leu Ile Asp Thr Asn Arg Ser Arg Ala Cys 180 185

His Pro Cys Ser Pro Met Cys Lys Gly Ser Arg Cys Trp Gly Glu Ser

195 20	0 205
Ser Glu Asp Cys Gln Ser Leu Th	r Arg Thr Val Cys Ala Gly Gly Cys
210 215	220
Ala Arg Cys Lys Gly Pro Leu Pro	o Thr Asp Cys Cys His Glu Gln Cys
225 230	235 240
Ala Ala Gly Cys Thr Gly Pro Lys	s His Ser Asp Cys Leu Ala Cys Leu
245	250 255
His Phe Asn His Ser Gly Ile Cys	Glu Leu His Cys Pro Ala Leu Val
260	265 270
Thr Tyr Asn Thr Asp Thr Phe Glu	Ser Met Pro Asn Pro Glu Gly Arg
275 280	285
Tyr Thr Phe Gly Ala Scr Cys Val	Thr Ala Cys Pro Tyr Ash Tyr Leu
290 295	300
Ser Thr Asp Val Gly Scr Cys Thr	Lou Val Cys Pro Lou His Asn Gln
305 310	315 320
Glu Val Thr Ala Glu Asp Gly Thr	Gln Arg Cys Glu Lys Cys Ser Lys
325	330 335
Pro Cys Ala Arg Val Cys Tyr Gly	Leu Gly Mot Glu His Leu Arg Glu
340	345 - 350
Val Arg Ala Val Thr Ser Ala Asn	Ile.Gln Glu Pho Ala Gly Cys Lys
355 360	365
Lys Ile Phe Gly Ser Leu Ala Phe I	Leu Pro Glu Ser Phe Asp Gly Asp
370 375	380
Pro Ala Ser Asn Thr Ala Pro Leu G	Eln Pro Glu Gln Leu Gln Val Phe
385 390	395 400
Glu Thr Leu Glu Glu Ile Thr Gly T	yr Leu Tyr Ile Ser Ala Trp Pro
405	410 415
Asp Ser Leu Pro Asp Leu Ser Val P. 420 4:	ne Gln Asn Leu Gln Val Ile Arg 25 430
Gly Arg Ile Leu His Asn Gly Ala Ty	yr Ser Leu Thr Leu Gln Gly Leu 445
Gly Ile Ser Trp Leu Cly Leu Arg Sc	r Leu Arg Glu Leu Gly Ser Gly

450	455	460	
Leu Ala Leu Ile F 465	His His Asn Thr His L 470	eu Cys Phe Val His Thr Val	
		is Gln Ala Leu Leu His Thi 90 495	-
Ala Asn Arg Pro G	lu Asp Glu Cys Val G	ly Glu Gly Leu Ala Cys His	3
500	505	510	
Gln Leu Cys Ala A	rg Gly His Cys Trp Gl	ly Pro Gly Pro Tnr Gln Cys	;
515	520	525	
Val Asn Cys Ser G	ln Phe Lew Arg Gly Gl	n Giu Cys Val Glu Glu Cys	
530	535	540	
Arg Val Leu Glm Gl	ly Leu Pro Arg Glu Ty	r Val Asn Ala Arg His Cys	
545	550	555 560	
Leu Pro Cys His Pr 56		n Asn Gly Ser Val Thr Cys 0 575	
Phe Gly Pro Glu Al.	a Asp Gln Cys Val Ala	a Cys Ala His Tyr Lys Asp	
580	585	590	
Pro Pro Phe Cys Va.	l Ala Arg Cys Pro Ser	Gly Val Lys Pro Asp Leu	
595	600	605	
Ser Tyr Met Pro Ile	e Trp Lys Phe Pro Asp	Glu Glu Gly Ala Cys Gln	
610	615	620	
Pro Cys Pro Ile Asn	Cys Thr His Ser Cys	Val Asp Leu Asp Asp Lys	
625	630	635 640	
Gly Cys Pro Ala Glu 645		Leu Thr Ser Ile Ile Ser 655	
Ala Val Val Gly Ile 660	Leu Leu Val Val Val	Leu Gly Val Val Phe Gly 670	
Ile Leu Ile Lys Arg	Arg Gln Gln Lys Ile	Arg Lys Tyr Thr Met Arg	
675	680	685	
Arg Leu Leu Glm Glu 690	Thr Glu Leu Val Glu 695	Pro Leu Thr Pro Ser Gly	
Ala Met Pro Asn Gln	Ala Gln Met Arg Ile 1	Leu Lys Glu Thr Glu Leu	

705	710	715	720
Arg Lys Val I	bys Val Leu Gly Ser 725	Gly Ala Phe Gly Th	nr Val Tyr Lys 735
-	le Pro Asp Gly Glu	Asn Val Lys Ile Pr	o Val Ala Ile
	40	745	750
Lys Val Leu A 755	rg Glu Asn Thr Ser 760	Pro Lys Ala Asn Ly	
Asp Glu Ala T	yr Val Met Ala Gly	Val Gly Ser Pro Ty	r Val Ser Arg
770	775	780	
Lou Leu Gly II	le Cys Lou Thr Sor	Thr Val Gln Leu Va.	l Thr Gln Leu
785	790	795	800
Met Pro Tyr Gl	y Cys Leu Leu Asp	His Val Arg Glu Asr	n Arg Gly Arg
	805	810	815
Leu Gly Ser Gl	-	Trp Cys Met Gln Ile	e Ala Lys Gly
82		825	830
Met Ser Tyr Le	u Glu Asp Val Arg 1	Leu Val His Arg Asp	
835	840	845	
Arg Asn Val Let	u Val Lys Ser Pro <i>F</i>	Asn His Val Lys Ile	Thr Asp Phe
850	855	860	
Gly Lou Ala Arg	g Leu Leu Asp Ile A	sp Glu Thr Glu Tyr	His Ala Asp
365	870	875	880
Gly Gly Lys Val	. Pro Ile Lys Trp M	et Ala Leu Glu Ser	Ile Leu Arg
	885	890	895
Arg Arg Phe Thr	· · · · · · · · · · · · · · · · · · ·	al Trp Ser Tyr Gly 05	Val Thr Val 910
Trp Glu Lou Met 915	Thr Phe Gly Ala Ly	ys Pro Tyr Asp Gly 925	Ile Pro Ala
Arg Glu Ile Pro	Asp Leu Leu Glu Ly	s Gly Clu Arg Leu	Pro Gln Pro
930	935	940	
Pro Ile Cys Thr	Ile Asp Val Tyr Me	t Ile Met Val Lys (Cys Trp Met
945		955	960
Ile Asp Ser Glu	Cys Arg Pro Arg Ph	e Arg Glu Leu Val S	Ser Glu Phe

Ser Arg Met Ala Arg Asp Pro Gln Arg Phe Val Val Ile Gln Asn Glu Asp Leu Gly Pro Ala Ser Pro Leu Asp Ser Thr Phe Tyr Arg Ser Leu Leu Glu Asp Asp Met Gly Asp Leu Val Asp Ala Glu Glu Tyr Leu 1015 1020 Val Pro Gln Gln Gly Phe Phe Cys Pro Asp Pro Ala Pro Gly Ala Gly Gly Mot Val His His Arg His Arg Ser Ser Ser Thr Arg Ser Gly Gly Gly Asp Lou Thr Leu Gly Lou Glu Pro Ser Glu Glu Glu Ala Pro Arg Ser Pro Leu Ala Pro Ser Glu Gly Ala Gly Ser Asp Val Phe Asp Gly Asp Leu Gly Met Gly Ala Ala Lys Gly Leu Gln Ser Leu Pro Thr His Asp Pro Ser Pro Lou Gln Arg Tyr Ser Glu Asp Pro Thr Val Pro Leu Pro Ser Glu Thr Asp Gly Tyr Val Aia Pro Leu Thr Cys Ser Pro Gln Pro Glu Tyr Val Asn Gln Pro Asp Val Arg Pro Gln Pro Pro Ser Pro 1145 1150 Arg Glu Cly Pro Leu Pro Ala Ala Arg Pro Ala Gly Ala Thr Leu Glu Arg Pro Lys Thr Leu Ser Pro Gly Lys Asn Gly Val Val Lys Asp Val Phe Ala Phe Gly Gly Ala Val Clu Ash Pro Glu Tyr Leu Thr Pro Gln Gly Gly Ala Ala Pro Gln Pro His Pro Pro Pro Ala Phe Ser Pro Ala

Phe Asp Ash Leu Tyr Tyr Trp Asp Gln Asp Pro Pro Glu Arg Gly Ala

1220 1225 1230

Pro Pro Ser Thr Phe Lys Gly Thr Pro Thr Ala Glu Asn Pro Glu Tyr 1235 1240 1245

Leu Gly Leu Asp Val Pro Val 1250 1255

<210> 6

<211> 1210

<212> PRT

<213> HUMAN

<220>

<223> numan EGFR

<400> 6

Met Arg Pro Ser Gly Thr Ala Gly Ala Ala Leu Leu Ala Leu Leu Ala 1 5 10 15

Ala Leu Cys Pro Ala Ser Arg Ala Leu Glu Glu Lys Lys Val Cys Gln
20 25 30

Gly Thr Ser Asn Lys Leu Thr Gln Leu Gly Thr Phe Glu Asp His Phe 35 40 45

Leu Ser Leu Gln Arg Met Phe Asn Asn Cys Glu Val Val Leu Gly Asn 50 55 60

Leu Glu Ile Thr Tyr Val Gln Arg Asn Tyr Asp Leu Ser Phe Leu Lys 65 70 75 80

Thr Ile Gln Glu Val Ala Gly Tyr Val Leu Ile Ala Leu Asn Thr Val 85 90 95

Glu Arg Ile Pro Leu Glu Asn Leu Gln Ile Ile Arg Gly Asn Met Tyr 100 105 110

Tyr Glu Asn Ser Tyr Ala Leu Ala Val Leu Ser Asn Tyr Asp Ala Asn 115 120 125

Lys Thr Gly Leu Lys Glu Leu Pro Met Arg Asn Leu Gln Glu Ile Leu 130 135 140

His Gly Ala Val Arg Phe Ser Asn Asn Pro Ala Leu Cys Asn Val Glu 145 150 155 160

Ser Ile Gin Trp Arg Asp Ile Val Ser Ser Asp Phe Leu Ser Asn Met 165 170 Ser Met Asp Phe Gln Asn His Leu Gly Ser Cys Gln Lys Cys Asp Pro 135 Ser Cys Pro Asn Gly Ser Cys Trp Gly Ala Gly Glu Asn Cys Gln 200 205 195 Lys Lou Thr Lys Ile Ile Cys Ala Gln Cln Cys Ser Gly Arg Cys Arg 210 215 Gly Lys Ser Pro Ser Asp Cys Cys His Asn Gln Cys Ala Ala Gly Cys 230 235 Thr Gly Pro Arg Glu Ser Asp Cys Leu Val Cys Arg Lys Phe Arg Asp 250 Glu Ala Thr Cys Lys Asp Thr Cys Pro Pro Leu Met Leu Tyr Asn Pro 260 265 Thr Thr Tyr Gln Met Asp Val Asn Pro Glu Gly Lys Tyr Sor Phe Gly 275 280 Ala Thr Cys Val Lys Lys Cys Pro Arg Asn Tyr Val Val Thr Asp His 300 290 295 Gly Ser Cys Val Arg Ala Cys Gly Ala Asp Ser Tyr Glu Met Glu Glu 305 310 315 Asp Cly Val Arg Lys Cys Lys Lys Cys Glu Gly Pro Cys Arg Lys Val 330 Cys Asn Gly Ile Gly Ile Gly Glu Phe Lys Asp Ser Leu Ser Ile Asn 340 345 Ala Thr Asn Ile Lys His Phe Lys Asn Cys Thr Ser Ile Ser Gly Asp 360 Leu His Ile Leu Pro Val Ala Phe Arg Gly Asp Ser Phe Thr His Thr 370 375 380 Pro Pro Leu Asp Pro Gln Glu Leu Asp Ile Leu Lys Thr Val Lys Glu 385 390 395 400

Ile Thr Gly Phe Leu Leu Ile Gln Ala Trp Pro Glu Asn Arg Thr Asp

410

405

Leu His Ala Phe Glu Ash Leu Glu lle Ile Arg Gly Arg Thr Lys Gln 425 His Gly Gln Phe Ser Leu Ala Val Val Ser Leu Asn Ile Thr Ser Leu 435 440 Gly Lou Arg Ser Lou Lys Glu Ile Ser Asp Gly Asp Val Ile Ile Ser 455 Cly Asn Lys Asn Leu Cys Tyr Ala Asn Thr Ile Asn Trp Lys Lys Leu 470 475 465 Phe Gly Thr Ser Gly Gln Lys Thr Lys Ile Ile Ser Asn Arg Gly Glu 490 485 Asn Ser Cys Lys Ala Thr Gly Gln Val Cys His Ala Leu Cys Ser Pro 500 505 Glu Gly Cys Trp Gly Pro Glu Pro Arg Asp Cys Val Ser Cys Arg Asn 515 520 Val Ser Arg Gly Arg Glu Cys Val Asp Lys Cys Lys Leu Leu Glu Gly 535 Glu Pro Arg Glu Phe Val Glu Asn Ser Glu Cys Ile Gln Cys His Pro 550 555 Glu Cys Leu Pro Gln Ala Met Asn Ile Thr Cys Thr Gly Arg Gly Pro 565 570 Asp Asn Cys Ile Gln Cys Ala His Tyr Ile Asp Gly Pro His Cys Val 580 585 Lys Thr Cys Pro Ala Gly Val Met Gly Glu Asn Asn Thr Leu Val Trp 595 600 Lys Tyr Ala Asp Ala Gly His Val Cys His Leu Cys His Pro Asn Cys 610 615 620 Thr Tyr Gly Cys Thr Gly Pro Gly Leu Glu Gly Cys Pro Thr Asn Gly 625 630 635 Pro Lys Ile Pro Ser Ile Ala Thr Gly Mct Val Gly Ala Leu Leu 650 Leu Leu Val Val Ala Leu Gly Ile Gly Leu Phe Met Arg Arg Arg His 660 665

Ile Val Arg Lys Arg Thr Leu Arg Arg Leu Leu Gln Glu Arg Glu Leu Val Glu Pro Lea Thr Pro Ser Gly Glu Ala Pro Ash Gln Ala Leu Leu Arg Ile Leu Lys Glu Thr Glu Phe Lys Lys Ile Lys Val Leu Gly Ser Gly Ala Phe Gly Thr Val Tyr Lys Gly Leu Trp Ile Pro Glu Gly Glu Lys Val Lys Ile Pro Val Ala Ile Lys Glu Leu Arg Glu Ala Thr Ser Pro Lys Ala Ash Lys Glu Ile Lou Asp Glu Ala Tyr Val Met Ala Ser Val Asp Asn Pro His Val Cys Arg Leu Leu Gly Ile Cys Leu Thr Ser Thr Val Gln Leu Ile Thr Cln Leu Met Pro Phe Gly Cys Leu Leu Asp Tyr Val Arg Glu His Lys Asp Asn Ile Gly Ser Gln Tyr Leu Leu Asn Trp Cys Val Glm Ile Ala Lys Gly Met Asn Tyr Leu Glu Asp Arg Arg Leu Val His Arg Asp Leu Ala Ala Arg Ash Val Leu Val Lys Thr Pro 835 840 945 Gln His Val Lys Ile Thr Asp Phe Gly Leu Ala Lys Leu Leu Gly Ala Glu Glu Lys Glu Tyr His Ala Glu Gly Gly Lys Val Pro Ile Lys Trp Met Ala Leu Gli Scr Ilc Leu His Arg Ilc Tyr Thr His Gln Ser Asp Val Trp Ser Tyr Gly Val Thr Val Trp Glu Leu Met Thr Phe Gly Ser Lys Pro Tyr Asp Gly Ile Pro Ala Ser Glu Ile Ser Ser Ile Leu Glu

Lys Gly Glu Arg Leu Pro Gln Pro Pro Fle Cys Thr Ile Asp Val Tyr 920 935 940

Met Ile Met Val Lys Cys Trp Met Ile Asp Ala Asp Ser Arg Pro Lys 945 950 955 960

Pho Arg Glu Leu Ile Ile Glu Phe Ser Lys Met Ala Arg Asp Pro Gln 965 970 975

Arg Tyr Leu Val Ile Gln Gly Asp Glu Arg Met His Leu Pro Ser Pro 980 985 990

Thr Asp Ser Ash Phe Tyr Arg Ala Leu Met Asp Glu Glu Asp Met Asp 995 1000 1005

Asp Val Val Asp Ala Asp Glu Tyr Lou Ile Pro Gln Gln Gly Phe Phe 1010 1015 1020

Ser Ser Pro Ser Thr Ser Arg Thr Pro Leu Leu Ser Ser Leu Ser Ala 1025 1030 1035 1040

Thr Ser Asn Asn Ser Thr Val Ala Cys Ilc Asp Arg Asn Gly Leu Gln
1045 1050 1055

Ser Cys Pro Ile Lys Glu Asp Ser Phe Leu Gln Arg Tyr Ser Ser Asp 1060 1065 1070

Pro Thr Gly Ala Leu Thr Glu Asp Ser Ile Asp Asp Thr Phe Leu Pro 1075 1080 1085

Val Pro Glu Tyr Ile Asn Gln Ser Val Pro Lys Arg Pro Ala Gly Ser 1090 1095 1100

Val Gln Asn Pro Val Tyr His Asn Gln Pro Leu Asn Pro Ala Pro Ser 1105 1110 1115 1120

Arg Asp Pro His Tyr Gln Asp Pro His Ser Thr Ala Val Gly Asn Pro 1125 1130 1135

Glu Tyr Leu Asn Thr Val Gln Pro Thr Cys Val Asn Ser Thr Phe Asp 1140 1145 1150

Ser Pro Ala His Trp Ala Gln Lys Gly Scr His Gln Ile Ser Leu Asp 1155 1160 1165

Ash Pro Asp Tyr Gln Gln Asp Phe Phe Pro Lys Glu Ala Lys Pro Ash 1176 1175 1180 Gly Ile Phe Lys Gly Ser Thr Ala Glu Asn Ala Glu Tyr Leu Arg Val 1185 1190 1195 1200

Ala Pro Gln Scr Scr Glu Phe Tle Gly Ala 1205 1210